



(877) 727-6800 www.antechimagingervices.com

Doctor's Copy

## PennHIP Report

Referring Veterinarian: Dr Wayne Griffin  
 Email: [REDACTED]

Clinic Name: Chippewa Veterinary Clinic  
 Clinic Address: 14961 81st Ave.  
 Chippewa Falls, WI 54729  
 Phone: (715) 723-3655  
 Fax: (715) 723-5245

## Patient Information

Client: Mars, Sandra  
 Patient Name: Leighla  
 Reg. Name: McJare's Flower Of The Night  
 Leighla  
 PennHIP Num: 109674  
 Species: Canine  
 Date of Birth: 29 Feb 2016  
 Sex: Female  
 Date of Study: 20 Jul 2017  
 Date of Report: 24 Jul 2017

Tattoo Num:  
 Patient ID: 57664  
 Registration Num: PR19033203  
 Microchip Num:  
 Breed: STANDARD POODLE  
 Age: 17 months  
 Weight: 43 lbs/19.5 kgs  
 Date Submitted: 24 Jul 2017

## Findings


Distraction Index (DI): Right DI = 0.43, Left DI = 0.52.  
 Osteoarthritis (OA): No radiographic evidence of OA for either hip.  
 Cavitation/Other Findings: None.

## Interpretation

Distraction Index (DI): The laxity ranking is based on the hip with the greater laxity (larger DI). In this case the DI used is 0.52.

OA Risk Category: The DI is between 0.50 and 0.69. This patient is at moderate risk for hip OA.

Distraction Index Chart:



**Breed Statistics:** This interpretation is based on a cross-section of 2910 canine patients of the STANDARD POODLE breed in the AIS PennHIP database. The gray strip represents the central 90% range of DIs (0.30 - 0.72) for the breed. The breed average DI is 0.48 (solid square). The patient DI is the solid circle (0.52).

**Summary:** The degree of laxity (DI = 0.52) falls within the central 90% range of DIs for the breed. This amount of hip laxity places the hip at a moderate risk to develop hip OA. No radiographic evidence of OA for either hip.

**Interpretation and Recommendations:** No OA/Moderate Risk: Likely to develop radiographic evidence of hip OA by 1-10 years of age (70% of dogs.) The risk to develop OA, the timing of OA onset, and the rate of progression are dependent upon many factors including DI, breed, body weight, age, and activity levels. **Recommendations:** Evidence-based

strategies to lower the risk of dogs getting OA or to treat those having OA fall into 5 modalities.\* For detailed information, consult these documents.\* Use any or all of these modalities as needed:

- 1) For acute or chronic pain prescribe NSAID PO short or long term. Amantadine can be added if response is marginal or if neuropathic pain is suspected.
- 2) Optimize body weight, keep lean, at BCS = 5/9.
- 3) Prescribe therapeutic exercise at intensities that do not precipitate lameness.
- 4) Administer polysulfated glycosaminoglycans IM or SQ, so-called DMOAD.
- 5) Feed an EPA-rich prescription diet preventatively for dogs at risk for OA or therapeutically for dogs already showing radiographic signs of OA.

At the present time there is inadequate evidence to confidently recommend any of the many other remedies to prevent or treat OA. Studies are in progress. Consider repeating radiographs at periodic intervals to determine the rate of OA progression and adjust treatment accordingly. Older dogs may show clinical signs such as chronic pain, reluctance to go stairs or jump onto the bed, and stiffness particularly after resting. It is unlikely that end-stage hip disease will develop for dogs at this risk level so surgical therapy for the pain of hip OA would rarely be indicated.

**Breeding Recommendations:** Please consult the PennHIP Manual.

\* From WSAVA Global Pain Council Guidelines and the 2015 AAHA/AAFP Pain Management Guidelines

**Comments:**

None